

for further human studies of limb ischaemia to induce a systemic preconditioned state.

Authors' affiliations

R K Kharbanda, University of Cambridge, Cambridge, UK
J Li, **I E Konstantinov**, **M M H Cheung**, **H Frndova**, **J Stokoe**, **P Cox**,
G Van Arsdell, **A N Redington**, The Hospital for Sick Children, Toronto,
 Ontario, Canada
P A White, Papworth Hospital, Cambridge, UK
M Vogel, **R MacAllister**, University College London, London, UK

This work was supported by a grant from the Canadian Institutes of Health Research. The British Heart Foundation supports RKK. MMHC is supported by the Heart and Stroke Foundation of Ontario, Canada

REFERENCES

- 1 **Chew MS**, Brandslund I, Brix-Christensen V, *et al*. Tissue injury and the inflammatory response to pediatric cardiac surgery with cardiopulmonary bypass: a descriptive study. *Anesthesiology* 2001;**94**:745–53.
- 2 **Massoudy P**, Zahler S, Becker BF, *et al*. Evidence for inflammatory responses of the lungs during coronary artery bypass grafting with cardiopulmonary bypass. *Chest* 2001;**119**:31–6.
- 3 **Yellon DM**, Alkhulaifi AM, Pugsley WB. Preconditioning the human myocardium. *Lancet* 1993;**342**:276–7.
- 4 **Kharbanda RK**, Mortensen UM, White PA, *et al*. Transient limb ischemia induces remote ischemic preconditioning in vivo. *Circulation* 2002;**106**:2881–3.
- 5 **Chaturvedi RR**, Hjortdal VE, Stenbog EV, *et al*. Inhibition of nitric oxide synthesis improves left ventricular contractility in neonatal pigs late after cardiopulmonary bypass. *Heart* 1999;**82**:740–4.
- 6 **Rimensberger PC**, Cox PN, Frndova H, *et al*. The open lung during small tidal volume ventilation: concepts of recruitment and "optimal" positive end-expiratory pressure. *Crit Care Med* 1999;**27**:1946–52.
- 7 **Li J**, Schulze-Neick I, Lincoln C, *et al*. Oxygen consumption after cardiopulmonary bypass surgery in children: determinants and implications. *J Thorac Cardiovasc Surg* 2000;**119**:525–33.
- 8 **Hoffman TM**, Wernovsky G, Atz AM, *et al*. Efficacy and safety of milrinone in preventing low cardiac output syndrome in infants and children after corrective surgery for congenital heart disease. *Circulation* 2003;**107**:996–1002.
- 9 **Perrault LP**, Menasche P, Bel A, *et al*. Ischemic preconditioning in cardiac surgery: a word of caution. *J Thorac Cardiovasc Surg* 1996;**112**:1378–86.
- 10 **Schulze-Neick I**, Li J, Penny DJ, *et al*. Pulmonary vascular resistance after cardiopulmonary bypass in infants: effect on postoperative recovery. *J Thorac Cardiovasc Surg* 2001;**121**:1033–9.
- 11 **Ng CS**, Wan S, Yim AP, *et al*. Pulmonary dysfunction after cardiac surgery. *Chest* 2002;**121**:1269–77.
- 12 **Massoudy P**, Zahler S, Becker BF, *et al*. Significant leukocyte and platelet retention during pulmonary passage after declamping of the aorta in CABG patients. *Eur J Med Res* 1999;**4**:178–82.
- 13 **Harkin DW**, Barros D'sa AA, McCallion K, *et al*. Ischemic preconditioning before lower limb ischemia-reperfusion protects against acute lung injury. *J Vasc Surg* 2002;**35**:1264–73.
- 14 **Charpie JR**, Dekeon MK, Goldberg CS, *et al*. Serial blood lactate measurements predict early outcome after neonatal repair or palliation for complex congenital heart disease. *J Thorac Cardiovasc Surg* 2000;**120**:73–80.
- 15 **Davies AR**, Bellomo R, Raman JS, *et al*. High lactate predicts the failure of intraaortic balloon pumping after cardiac surgery. *Ann Thorac Surg* 2001;**71**:1415–20.
- 16 **Przklenk K**, Darling CE, Dickson EW, *et al*. Cardioprotection 'outside the box'. *Basic Res Cardiol* 2003;**98**:149–57.
- 17 **Konstantinov IE**, Arab S, Kharbanda RK, *et al*. The remote ischemic preconditioning stimulus modifies inflammatory gene expression in humans. *Physiol Genomics* 2004;**19**:143–50.


IMAGES IN CARDIOLOGY

doi: 10.1136/hrt.2005.087098

Malignant pectus excavatum

A 7-year-old girl with anterior Raphe syndrome was seen for recurrent syncope (six episodes over six months). She had undergone sternal repair of a midline defect at 3 years of age. The majority of episodes occurred in the morning and were usually associated with showering. Of concern was that an episode had occurred while swimming, requiring her to be rescued from the pool. Excepting obvious pectus excavatum, her clinical examination was normal. Her ECG, 24 ambulatory ECG and exercise test were normal and there was no evidence of long QT syndrome. Her transthoracic echocardiogram showed a structurally and functionally normal heart but the sternum appeared to compress the right ventricle. There was no flow acceleration across the inflow of the right ventricle. Syncope continued despite fluid and salt loading. A chest computed tomographic (CT) scan showed significant right ventricular compression by the sternal deformity (see panel). In the absence of any other cause for her symptoms we undertook sternochondroplasty with a resulting notable improvement in her pectus excavatum. She remains well and has had no further episodes of syncope at two years of follow up from her surgery.

Pectus excavatum is usually a cosmetic deformity but symptomatic cases of compression of the thoracic vessels and heart are described. To our knowledge this is a unique case presenting with syncope, which was presumably due to the flow limitation caused by the fixed right ventricular compression.



Chest computed tomographic scan with contrast showing right ventricular (RV) compression by the pectus excavatum.

D S Crossland
A W Auld
A M Davis
 davidxland@hotmail.com